

Historic, Archive Document

Do not assume content reflects current scientific knowledge, policies, or practices.

aQC981
.8
.C5G652

.8

.C5G652

RECEIVED
FEB 11 1964

16.00

ORDS
144-150
151-155
156-160
161-165
166-170
171-175
176-180
181-185
186-190
191-195
196-200
201-205
206-210
211-215
216-220
221-225
226-230
231-235
236-240
241-245
246-250
251-255
256-260
261-265
266-270
271-275
276-280
281-285
286-290
291-295
296-300
301-305
306-310
311-315
316-320
321-325
326-330
331-335
336-340
341-345
346-350
351-355
356-360
361-365
366-370
371-375
376-380
381-385
386-390
391-395
396-400
401-405
406-410
411-415
416-420
421-425
426-430
431-435
436-440
441-445
446-450
451-455
456-460
461-465
466-470
471-475
476-480
481-485
486-490
491-495
496-500
501-505
506-510
511-515
516-520
521-525
526-530
531-535
536-540
541-545
546-550
551-555
556-560
561-565
566-570
571-575
576-580
581-585
586-590
591-595
596-600
601-605
606-610
611-615
616-620
621-625
626-630
631-635
636-640
641-645
646-650
651-655
656-660
661-665
666-670
671-675
676-680
681-685
686-690
691-695
696-700
701-705
706-710
711-715
716-720
721-725
726-730
731-735
736-740
741-745
746-750
751-755
756-760
761-765
766-770
771-775
776-780
781-785
786-790
791-795
796-800
801-805
806-810
811-815
816-820
821-825
826-830
831-835
836-840
841-845
846-850
851-855
856-860
861-865
866-870
871-875
876-880
881-885
886-890
891-895
896-900
901-905
906-910
911-915
916-920
921-925
926-930
931-935
936-940
941-945
946-950
951-955
956-960
961-965
966-970
971-975
976-980
981-985
986-990
991-995
996-1000

Natural Resources Institute

Global Climate Change Research

1991 Status Report

March 1992



*U.S. Department of Agriculture
Agricultural Research Service
Beltsville Agricultural Research Center*

Beltsville Agricultural Research Center

Natural Resources Institute

Global Climate Change Research

Status Report - 1991

March 1992

Compiled by:

*Donald T. Krizek
Climate Stress Laboratory
Natural Resources Institute*

A COMPILATION OF GLOBAL CLIMATE CHANGE RESEARCH AT THE
BELTSVILLE AGRICULTURAL RESEARCH CENTER CY 1991
Donald T. Krizek

INTRODUCTION

This report covers the activities of scientists at the Beltsville Agricultural Research Center (BARC) involved in research on global climate change research. The report covers work done primarily in four laboratories in the Natural Resources Institute in the areas of hydrology, remote sensing, systems research, and climate stress effects. Work conducted in the Plant Science Institute in the Molecular Biology Laboratory and Weed Science Laboratory is also included.

SECTION 1.

This section covers highlights of research conducted in CY 1991. A short description is given of the purpose, accomplishment, and significance of the research. For further information, the reader is referred to the Annual CRIS Reports.

SECTION 2.

This section contains a list of peer-reviewed papers published during 1989 to 1991.

SECTION 3.

This section contains a list of abstracts of papers published during 1989 to 1991.

SECTION 4.

This section contains a selected list of papers submitted for ARS approval between January 1990 and December 1991 followed by an interpretive summary and abstract of the manuscript. This section was obtained through the ARS TEXTRAN (Technology Transfer Automated Retrieval System) and is available upon request.

SUMMARY

BARC scientists are actively involved in research to help better understand possible physical and biological changes that might result from global climate change. Research is underway: 1) to develop improved data bases and models of plant growth; 2) to determine possible hydrologic changes; 3) to develop improved radiative transfer models; 4) to understand the biochemical and physiological mechanisms of adaptation to CO₂ enriched atmospheres, air pollutants, and increased UV-B radiation; and 5) to determine the interactive effects of CO₂ and other environmental factors including tropospheric ozone, UV-B radiation, and water stress.

DISCLAIMER

The author accepts full responsibility for any errors of omission or commission that have been made. Persons inadvertently omitted from inclusion from this compilation are encouraged to contact me by phone (301-504-5324) or by fax (301-504-7521). Technical questions concerning specific areas of research should be directed to the contact scientist listed.

SECTION 1

RESEARCH ACCOMPLISHMENTS

1991

1991 ANNUAL REPORT ON GLOBAL CLIMATE CHANGE RESEARCH AT BARC

A. HYDROLOGIC STUDIES

MONSOON '90 - AN INTERDISCIPLINARY FIELD EXPERIMENT FOR QUANTIFYING BASIN SCALE HYDROLOGIC FLUXES IN SEMIARID REGIONS

An interdisciplinary field experiment in evaluating the water and energy fluxes of a semi-arid watershed was conducted in Arizona over the summer of 1990. Ground truth and remote sensing measurements covered a wide range of spatial and temporal scales. This data is being utilized in energy and water balance models which employ remote sensing to improve our capabilities of integrating hydrologic fluxes to the basin scale. The findings can then be incorporated in large scale atmospheric models simulating the interaction of hydrologic processes with the atmosphere for semiarid climates. This will ultimately yield more reliable modeling of the impacts of climate change on the hydrologic cycle over different ecosystems.

CRIS 1270-13000-004-00D

Dr. W. Kustas

SNOWMELT MODEL TESTED WORLDWIDE IN DIFFERENT CLIMATES

The Snowmelt-Runoff Model (SRM) has been tested around the world in different climates to establish baselines for use in comparisons of various climate change scenarios. SRM, a simple degree-day model with physically derived parameters, has been applied to over 50 basins in 15 different countries. SRM performance has been uniformly high for existing conditions (average $R^2 = 0.84$ and average volume difference = 3.8%) in all climate regions tested, and when consistent climate change scenarios are available in these regions, the model can be utilized to estimate a range of hydrologic responses that will allow formulation of possible management strategies.

CRIS 1270-13000-004-00D

Dr. A. Rango

HYDROLOGIC RESPONSE TO POTENTIAL CLIMATE CHANGE IN WESTERN NORTH AMERICAN BASINS

Projected climate changes in mountain basins of western North America will significantly influence snow accumulation, snowmelt, and subsequent water yield. The Snowmelt-Runoff Model (SRM) has been used on the Rio Grande (Colorado), Kings River (California), and Illecillewaet River (British Columbia) with various climate change scenarios. Results indicate that in wet years floods will become more numerous and in dry years drought will be more frequent. A redistribution of flow will also take place so that there will most likely be a widening of the gap between water supply and water demand in most river basins.

CRIS 1270-13000-004-00D

Dr. A. Rango

B. STUDIES ON REMOTE SENSING

REMOTE SENSING OF ABSORBED RADIATION AND PHYTOMASS PRODUCTION

Linear relations between phytomass production and absorbed photosynthetically active radiation (PAR) have been reported for a wide range of plant species. Other work has shown that absorbed PAR may be estimated from remotely sensed observations. The combination of these two concepts provides the basis of a physically-based agronomic monitoring system. For both corn and soybean, absorbed PAR was linearly related to remotely sensed reflectance measurements. The rate of accumulation of phytomass per unit absorbed PAR was greater for corn than for soybean. These results indicate that remotely sensed measurements contribute valuable information concerning energy and mass accumulations in plant canopies.

CRIS 1270-66000-010-00D

Dr. C. S. T. Daughtry

C. SYSTEMS RESEARCH

A TWO-DIMENSIONAL SIMULATOR OF PLANT GROWTH, AND WATER AND SOLUTE TRANSPORT

Crop simulation models often suffer in their predictive capabilities because they often have very simple representations of soil process. The Systems Research Laboratory scientists have coupled process-level crop simulation models with the two-dimensional, finite element model SOILSIM. This will greatly improve the modeling of plant growth and solute movement under a range of climatic conditions. The soybean model GLYCIM has been interfaced with SOILSIM and the Cotton Production Model (CPM) will incorporate SOILSIM.

CRIS 1270-66000-012-00D

Dr. D. Timlin

ESTIMATING EFFECTS OF CLIMATE CHANGE ON SOYBEAN PRODUCTION

In order for U.S. agriculture to adequately respond to global climate changes, a sound understanding of their effects on major agronomic crops is necessary. Effects of temperature and CO₂ elevation on soybeans are documented in the literature. However, the implications for regional production have not been investigated. The current work seeks to extend the use of simulation modelling of soybean growth to estimate the effects of global climate change on regional soybean yields. Using the simulation model GLYCIM in combination with the yield sampling methodology of the National Agricultural Statistics Service, USDA, yield estimates under future climatological conditions will be obtained. This research is vital to assess the effects of global climate change on this important crop so that adequate policy responses can be created and implemented.

CRIS 1270-61000-012-00D

Dr. J. Haskett

CARBON DIOXIDE AND TEMPERATURE EFFECTS ON COTTON

The effects of carbon dioxide concentration [CO_2], temperature, and their interactions were evaluated on cotton physiological processes such as growth, development, root growth, photosynthesis, and respiration. The study was conducted using controlled environmental cabinets with five temperature treatments of 15/7°C, 20/12°C, 25/17°C, 30/22°C, and 35/27°C (day/night) and CO_2 treatments of 350 and 700 $\mu\text{mol mol}^{-1}$ at each temperature. Plant height and the rate of stem elongation increased with increases in temperature and CO_2 concentration. The number of main stem nodes and the node addition rate are temperature dependent. In addition, CO_2 has no effect on these processes. The developmental processes such as squaring, flowering, and boll opening were driven by temperature. Photosynthesis increased linearly with increases in CO_2 concentration. This database is critical for the development of process-level crop simulation models which can be used for predicting the effects of future climate change on agriculture.

CRIS 1270-6100-012-00D Dr. V. R. Reddy

D. BIOCHEMICAL AND PHYSIOLOGICAL EFFECTS

IMPACT OF DROUGHT STRESS ON COTTON SEED AND FIBER PROPERTIES

Global change may result in shifting patterns of seasonal temperature and rainfall. The effect of severe drought (water withholding) on developing seeds in bolls of mature cotton plants grown under greenhouse conditions was investigated, using two cultivars of Gossypium hirsutum - Deltapine M-8 and DPL 51. Analyses of fiber and seed properties were made on mature seed from bolls on upper and lower branches of plants receiving 10 to 30 days drought stress. Data for fiber properties of seed (M-8) from bolls receiving 25 to 30 days drought stress provided evidence that these fibers were immature and took up little stain following a standard dye test. Radiolabeling studies with DPL 51 for incorporation into protein revealed a stage specific sensitivity to drought for continued incorporation into storage proteins. These findings will be used for development of a module for predicting changes in seed and fiber quality due to drought stress for field-grown cotton.

CRIS 1270-11210-004-00D

Dr. E. L. Vigil

ELEVATED CARBON DIOXIDE CONCENTRATION REDUCES RESPIRATION

Elevated atmospheric carbon dioxide concentrations increase photosynthesis, and it was thought they would also increase respiration. However, when respiration rates were examined in ten species grown at twice the current atmospheric carbon dioxide concentration, it was found that respiration rates were reduced at elevated carbon dioxide in nine of the species. Interactions between carbon dioxide concentration and temperature were examined in alfalfa and orchard grass leaves. Results indicate that the sensitivity of respiration to temperature declined with time in leaves grown at elevated carbon dioxide. This resulted in a larger effect of carbon dioxide on respiration as temperature increased. Inhibition of respiration by elevated carbon dioxide has potentially important consequences to crop carbon balance and to the future trend of atmospheric carbon dioxide concentrations.

CRIS 1270-21000-011-00D

Drs. J. A. Bunce and L. H. Ziska

RAPID PHOTOSYNTHETIC ADJUSTMENT TO ELEVATED CARBON DIOXIDE

Elevated atmospheric carbon dioxide usually stimulates photosynthesis initially, but often physiological adjustments occur which decrease or eliminate the stimulation. Experiments with soybeans and sugarbeets used variation in light, temperature and nutrients to test the hypotheses that photosynthetic adjustment to elevated carbon dioxide resulted from feedback inhibition or nutrient deficiency. The patterns of response were not consistent with either hypothesis. In soybeans, a single dark period at 700 ppm carbon dioxide decreased photosynthetic rates the next day compared to control leaves kept at 350 ppm. This suggests a regulatory response, rather than a change in leaf composition was involved. In experiments with orchard grass and alfalfa, it was found that photosynthetic adjustment to elevated carbon dioxide increased with increased growth temperature in alfalfa, but decreased in orchard grass. The patterns of photosynthetic acclimation were consistent with a smaller effect of elevated carbon dioxide on growth rate at warmer temperatures in alfalfa, but a larger growth stimulation at warmer temperatures in orchard grass. A better understanding of these responses will be necessary in order to predict crop growth at elevated atmospheric carbon dioxide concentrations.

CRIS 1270-21000-011-00D

Drs. J. A. Bunce and L. H. Ziska

ANTICIPATED INCREASES IN TEMPERATURE AND CO₂ MAY AFFECT SEED GERMINATION

We examined the influence of 35 and 70 Pa carbon dioxide pressure on germination in pea, corn, sunflower, soybean, amaranth, lambsquarters and alfalfa grown in controlled environment chambers at 20 and 30°C. Earlier seed germination (24-48 h) was noted at 70 Pa CO₂ for pea, corn, lambsquarters, and alfalfa. A significant increase in the total number of seeds germinated was also observed in lambsquarters and alfalfa with elevated CO₂. With the exception of amaranth, a 10°C rise in temperature did not influence the germination response to CO₂, although seeds overall germinated faster at the higher temperature. For amaranth, no significant germination was observed at 20°C, but earlier germination and a greater total germination percentage was noted at 30°C and high CO₂. Diurnal studies of seed imbibition showed a greater rate of water uptake for those seeds which responded to elevated CO₂. Results from this study suggest that changes in seed germination with anticipated increases in CO₂ and temperature may potentially alter stand establishment, crop cover and weed-crop interactions.

CRIS 1270-21000-011-00D

Dr. L.H. Ziska

STARCH SYNTHESIS IN THE LEAVES OF CO₂ ENRICHED SOYBEAN PLANTS

An examination of the response of soybean leaf starch synthesis to increases in atmospheric CO₂ concentration was undertaken. Glycine max [L.] Merr. cv Amsoy plants were propagated in growth chambers maintained with 650 $\mu\text{mol m}^{-2} \text{s}^{-1}$ white light, 14 h light-10 h dark cycle, 27°C continuous, and 65% RH, for 20 days post-emergence (PE) in an ambient atmosphere containing 330 $\mu\text{mol mol}^{-1}$ CO₂ air (normal CO₂ plants). At 20 days PE, one-half of the plants were transferred to a growth chamber with identical conditions except that the atmosphere in the chamber contained CO₂ at 1000 $\mu\text{mol mol}^{-1}$ air for 7 additional days (high CO₂ plants). After 7 days of high or normal CO₂ exposure, rates of foliar net photosynthesis (APS) were measured. Plant organs were sampled into liquid N₂, lyophilized, extracted, and quantitated for starch, sucrose, and phosphorylated metabolites. APS rates for source leaves of high CO₂ plants (measured in 1000 $\mu\text{mol mol}^{-1}$ CO₂ air) compared with those of normal CO₂ plants (in 330 $\mu\text{mol mol}^{-1}$ CO₂ air) were respectively, 1136 and 683 $\mu\text{mol CO}_2 \text{ fixed dm}^{-2} \text{ h}^{-1}$. During illumination, high compared with normal CO₂ plants exhibited 2-7 times higher starch levels in all organs including leaves, stems, and roots. Leaves from high compared with normal CO₂ plants possessed, respectively, $\approx 50\%$ and $\approx 7\%$ of their dry weight as starch. In contrast, relative to normal CO₂ plants, foliar sucrose levels in high CO₂ plants only were 1.3 to 1.6 times higher in the leaves and only slightly higher or not different in the stems and roots. The foliar concentrations of photosynthetic intermediates, e.g. glycerate-3-P (PGA), fructose-6-P (F6P) glucose-6-P (G6P), and glucose-1-P (G1P), were, in the high CO₂ plants, approximately twice those in CO₂ normal plants. Thus higher foliar levels of PGA, F6P, G6P and G1P in the high CO₂ plants suggested that a causal factor of elevated starch accumulation was higher activity of one or more rate limiting enzymes in the starch synthesis pathway; e.g. PGA is a positive effector and G1P is a substrate for ADP Glucose pyrophosphorylase, a very rate limiting enzyme in the starch synthesis pathway. The results and interpretations of this study contribute to a continuing effort to understand the physiological responses of soybean plants as they grow in a global atmosphere where the concentration of CO₂ is increasing.

CRIS 1270-21000-014-00D

Dr. J. Michael Robinson

INTERACTIONS BETWEEN AIR POLLUTANTS AND ATMOSPHERIC CO₂

Open-top field chambers and remote sensing techniques were used to investigate the leaf spectral response of winter wheat, corn, and soybean grown under enriched CO₂ concentrations and O₃ or CO₂ x SO₂ stress conditions. Photosynthesis and canopy leaf area index reflectance measurements were taken to evaluate the potential of remote sensing data to estimate losses in crop productivity due to air pollution. Laboratory studies have also been conducted on snapbean and soybean in environmentally controlled fumigation chambers. Determination of polyamine levels and PS II chlorophyll fluorescence of leaf tissues in response to antioxidant treatments and O₃ stress were also conducted. Effects of CO₂ and O₃ individually were consistent with those reported previously. The CO₂ and O₃ interactions observed were generally consistent with a reduction of O₃ injury by increased CO₂. The significance of these studies was to incorporate remote sensing technology to detect and assess the impact of atmospheric gases as it related to global climate changes. We would expect the research to be utilized not only in the scientific community and government but to agricultural chemical companies as well.

CRIS 1270-21000-008-00D

Dr. E. H. Lee

TEMPERATURE CROSS PROTECTION AGAINST SO₂ INJURY

The effects of temperature cross protection were investigated in two cultivars of coleus, differing in SO₂ sensitivity. Measurements were made of membrane lipid composition after 5 days of temperature pretreatment at 13 or 20C. Lipid composition at 20C revealed inherent differences between SO₂ sensitive 'Buckley Supreme' and SO₂ insensitive 'Marty'. Temperature hardening at 13C, which increased SO₂ tolerance, caused specific changes in membrane lipids. Determination of leaf free fatty acid pools and measurements of lipid peroxidation products further indicated a difference in metabolism between the two cultivars. These findings may help to elucidate the biochemical basis for SO₂ tolerance in plants.

CRIS 1275-24000-078-00D

Dr. H. A. Norman

CANOPY ARCHITECTURE, CO₂ AND SPECTRAL QUALITY

Elevated atmospheric CO₂ may stimulate vegetative growth, altering mutual shading of adjacent plants in densely planted monocultures. Canopy shade reduces both photosynthetically active radiation and the ratio of red:far-red radiation, affecting the photochromic pigment phytochrome. Many types of controlled environment lighting provide little far-red radiation, thus circumventing normal changes in spectral quality and altering phytochrome photostationary state. Soybean dry matter accumulation was less inhibited by high density planting when daylight red:far-red ratios were simulated in growth chambers. Improved growth was associated with reduced axillary development and increased stem and petiole extension. As a result, light penetration into the canopy improved and canopy photosynthesis increased. Morphological changes probably resulted from changes in the photostationary state of phytochrome. The influence of elevated CO₂ will be investigated. These results should be of interest to physiologists conducting experiments with elevated CO₂ in controlled environments and to breeders searching for strategies to improve growth and yield in response to rising CO₂.
CRIS 1270-11210-002-00D Dr. S. J. Britz

ADVERSE EFFECTS OF AMBIENT UV-B RADIATION LEVELS ON CROP GROWTH

UV-B radiation exclusion studies were conducted under field conditions in New Delhi, India during July, 1990, using a polyester filter placed horizontally over the plot to exclude the direct component of solar UV-B radiation. This filter excluded most of the direct UV radiation in the 280-320 nm region and transmitted only 23 percent of the UV radiation at 320 nm. The visible radiation at ground level was about 86% of the unfiltered radiation. In general, mungbean plants grown under the UV-B filter grew faster, had longer internodes, and greater leaf size than those exposed to full sunlight. There was a significant increase in pod and seed number, and dry weight of pods and seed when the direct component of UV-B radiation was excluded. These preliminary findings suggest that plants under natural atmospheric conditions in New Delhi are already growing under above optimal conditions of UV-B radiation. These findings should be of interest to researchers and policy makers involved in global climate change assessments.
CRIS 8001-11210-758-00P Drs. Y.P. Abrol, A.Sharma, & U.K.Sengupta;
Dr. D. T. Krizek, Cooperating Scientist

PHOTOPROTECTION MECHANISMS AGAINST DAMAGE FROM UV-B RADIATION

Sensitivity to ultraviolet-B radiation (UV-B, 280-315 nm) is generally reduced when background photosynthetically-active radiation (PAR) is high. The involvement of photosynthesis in the amelioration of UV-B damage was tested during short-term treatments at high PAR with high levels of supplemental UV-B and either "low" (450 vpm) or "high" (750 vpm) atmospheric CO₂. UV-B had little effect on cucumber leaf photosynthesis at low CO₂ but it inhibited leaf area expansion and plant growth. High CO₂ counteracted the effect of UV-B. In contrast, the accumulation of UV-absorbing flavonoid compounds was stimulated by UV-B exposure but was not affected by CO₂ enrichment. This result is consistent with the involvement of non-photosynthetic photoreceptors in the regulation of response to UV-B and should be of interest to physiologists seeking to assess the influence of elevated UV-B and breeders looking for traits that confer resistance to UV-B.

CRIS 1270-11210-002-01T Dr. S. J. Britz

INFLUENCE OF UV-B RADIATION ON POLYAMINES, LIPID PEROXIDATION & MEMBRANE LIPIDS

Cultivar differences in UV-B sensitivity in cucumber were first demonstrated at Beltsville in the mid-1970's. Current effort is being focused on establishing the biochemical basis for these genotype differences. UV-B sensitive (Poinsett) and UV-B insensitive (Ashley) cucumber were grown in the greenhouse for four weeks under 0 or 12.2 kJ m⁻² d⁻¹ of biologically effective UV-B radiation to determine effects of increased UV-B level corresponding to approximately a 20% depletion in stratospheric ozone level at Beltsville on June 21. Elevated UV-B radiation caused an increase in concentration of the diamine putrescine and the polyamine spermidine in cotyledon and leaf tissues in response in both cultivars, but levels were not correlated with UV-B sensitivity. Lipid peroxidation, as quantified by measurement of malondialdehyde (MDA) was increased by UV-B exposure. Significant cultivar differences were observed, with increases in lipid peroxidation in both cotyledons and leaves being correlated with UV-B sensitivity. Determination of membrane lipid composition indicated slight decreases in the unsaturated/saturated fatty acid ratios as a result of UV-B exposure. In general, Poinsett had a lower unsaturated/saturated ratio of fatty acids than did Ashley. These results indicate that membrane lipids may be a target of UV-B damage and that differences in UV-B susceptibility of these two cultivars may involve differences in lipid metabolism.

1270-11210-003-00D Drs. G. F. Kramer, H. A. Norman & D. T. Krizek

UV-B RADIATION MODIFIES PLANT RESPONSE TO ELEVATED TEMPERATURES

The effect of ultraviolet-B (UV-B, 280-315 nm) radiation on the thermal sensitivity of cucumber was examined to assess the potential agronomic consequence of combined stratospheric ozone depletion and global warming. Using an enzymatic assay (involving triphenyl tetrazolium chloride) to assess tissue viability after temperature (50C) treatment, it was apparent that growth of cucumber under enhanced UV-B radiation levels increased their ability to withstand elevated temperatures. The results suggest that both enhanced UV-B radiation and elevated temperatures may result in similar cellular lesions. Therefore, plant adaptation to elevated UV-B radiation results in protection against high temperatures.

CRIS 1270-21000-001-00T

Drs. C. R. Caldwell and A. Upadhyaha

ENHANCED TURNOVER OF THE D1 AND D2 PHOTOSYSTEM II REACTION CENTER PROTEINS UNDER UV-B IRRADIATION

The D1 and D2 heterodimer comprises the nucleus of the photosystem II (PSII) reaction center. D1 undergoes rapid degradation at low intensities of both visible and UV radiation, turning over at rates far exceeding those for other PSII proteins. The D2 protein degrades relatively slowly in visible light but in the UV its degradation rate increases considerably, approaching that of the D1 protein. Other PSII proteins are not as markedly affected. When a small amount of UV irradiation is added to saturating amounts of the visible light (as in sunlight) a synergistic stimulation of D1 degradation occurs, but even stronger stimulation of the D2 protein degradation is seen. Thus, a specific stimulation of the D2 protein degradation occurs in UV radiation, resulting in the D1-D2 heterodimer being the target of this damaging irradiation.

CRIS: 1275-21000-060-00D

Dr. A. K. Mattoo

METABOLIC CHANGES IN BEAN PLANTS UNDER UV- AND BLUE-DEFICIENT ENVIRONMENTS

'Bush Blue Lake' snapbean plants, known to be sensitive to ozone pollution and other oxidant stresses were grown for 4 weeks in growth chambers in the presence or absence of UV and blue radiation to investigate the biochemical basis for bleaching observed previously under UV and blue-deficient low pressure sodium (LPS) lamps. Plants were grown at $300 \mu\text{mol m}^{-2} \text{s}^{-1}$ of photosynthetically active radiation (PAR) under: a) LPS lamps, b) a combination of LPS and supplemental blue (B) fluorescent lamps; or c) under cool white fluorescent (CWF) lamps. By 20 days under LPS lamps, the compound leaves and leaves of the lateral shoots had developed extensive chlorosis, crinkling of the leaf margins, and severe interveinal necrosis. Exposure of plants to LPS + B lamps greatly ameliorated these symptoms, while plants grown under CWF lamps were free of injury. Electrophoretic separation of leaf proteins from these plants revealed significant differences in protein profiles under the three radiation sources. These findings provide further biochemical evidence for the essentiality of blue/UV radiation in normal chloroplast development.

CRIS 1270-11210-003-00D

A. L. Fleming

SECTION 2

PEER-REVIEWED PUBLICATIONS

1989 - 1991

1991 ANNUAL REPORT ON GLOBAL CLIMATE CHANGE RESEARCH AT BARC

A. HYDROLOGIC STUDIES

Engman, E. T., G. Angus, and W. P. Kustas. 1989. Relationships between the hydrologic balance of a small watershed and remotely-sensed soil moisture. IAHS-AISH, Pub. No. 186, pp. 75-84.

Engman, E. T., W. P. Kustas, and J. R. Wang. 1989. Remotely sensed soil moisture input to a hydrological model. Proceedings of the 1989 International Geosciences and Remote Sensing Symposium (IGARRS) Vancouver, Canada. 4:2150-2153.

Hipps, L. E., E. Swiatek, and W. P. Kustas. 1991. Effects of spatial variations in surface water balance on the surface atmospheric boundary layer interactions in an arid region. Preprint Vol. of the Tenth Conference. pp. 155-157.

Kustas, W. P. and C. S. T. Daughtry. 1990. Estimation of soil heat flux/net radiation ratio from spectral data. Agric. Forest Meteorol. 49:205-223.

Kustas, W. P., D. C. Goodrich, and M. S. Moran. 1991. An interdisciplinary field study of the energy and water fluxes in the atmosphere-biosphere system over semiarid rangelands: Description and some preliminary results. Bull. Amer. Meteorol. Soc. 72(11):1683-1700.

Kustas, W. P., M. S. Moran, R. D. Jackson, L. W. Gay, L. F. W. Duell, K. E. Kunkel, and A. D. Matthias. 1990. Instantaneous and daily values of the surface energy balance over agricultural fields using remote sensing and a reference field in an arid environment. Remote Sens. Environ. 32:125-141.

Kustas, W. P., M. S. Moran, P. J. Pinker, L. E. Hipps, E. Swiatek, D. I. Stannard, J. H. Blanford. 1991. Issues in mapping evapotranspiration over large areas with remote sensing data from MONSOON '90. Preprint Vol. of the Tenth Conference on Biometeorology and Aerobiology and Special Session on Hydrometeorology, 10-13 Sept. 1991, Salt Lake City, Utah. pp. 151-154.

Kustas, W. P., E. M. Perry, and P. C. Doraiswamy. 1991. MONSOON '90 - A remote sensing feasibility study of the energy and water balance of a semiarid basin. Preprint IAF-91-160 of the 42nd Congress of the International Astronautical Federation, 5-11 Oct. 1991, Montrael, Canada. p 6.

Martinec, J. and A. Rango. 1989. Effects of climate change on snowmelt runoff patterns, Remote Sensing and Large-Scale Global Processes, IAHS-AISH Pub. No. 186, International Association of Hydrological Sciences, pp. 31-38.

Moran, M. S., W. P. Kustas, L. B. Bach, M. A. Weltz, A. R. Huete, S. A. Amer. 1991. Use of remotely sensed spectral data for evaluation of hydrologic parameters in semiarid rangeland. Preprint Volume of the Tenth Conference on Biometerology. pp. 238-241.

Moran, M. S., W. P. Kustas, A. Vidal, D. I. Stannard, and J. H. Blanford. 1991. Use of ground-based remotely sensed data for surface energy balance calculations during MONSOON 90. Proceedings of the International Geoscience and Remote Sensing Symposium, Espoo, Finland, 3-6 June 1991, pp. 33-37.

Rango, A. 1991. Worldwide testing of the Snowmelt Runoff Model with applications for predicting the effects of climate change, *Nordic Hydrology* 22:10. (Abstract)

Rango, A. and V. van Katwijk. 1990. Climate change effects on the snowmelt hydrology of western North American mountain basins, *IEEE Transactions on Geoscience and Remote Sensing*, GE-28(5), pp. 970-974.

Rango, A. and V. van Katwijk. 1990. Water supply implications of climate change in western North American basins. Proceedings of the Symposium on International & Transboundary Water Resources Issues, American Water Resources Association, Toronto, Ontario, pp. 577-586.

Rango, A., J. Thurman, W. Blackburn, F. Schiebe, and D. Woolhiser. 1991. Compilation and availability of Agricultural Research Service hydrological data for use in climate change studies. Proceedings of the NATO Workshop on Opportunities for Hydrological Data in Support of Climate Change Studies, Lahnstein, Germany, NATO Publications Series.

van Katwijk, V. and A. Rango. 1991. The effect of climate change on snowmelt-runoff in two different climatic regions of the western United States, Proceedings of the American Meteorological Society Special Session on Hydrometeorology, Salt Lake City, Utah, pp. 210-213.

B. STUDIES ON REMOTE SENSING

Daughtry, C. S. T. 1990. Direct measurements of canopy structure. *Remote Sensing Reviews*. 5:45-60.

Daughtry, C. S. T., W. P. Kustas, M. S. Moran, P. J. Pinter, Jr., R. D. Jackson, P. W. Brown, W. D. Nichols, and L. W. Gay. 1990. Spectral estimates of net radiation and soil heat flux. *Remote Sensing Environ.* 32:111-124.

Kustas, W. P. and C. S. T. Daughtry. 1990. Estimation of the soil heat flux/net radiation ratio from spectral data. *Agricultural and Forest Meteorology* 49:205-223.

Prince, S. D., S. N. Goward, D. G. Dye, and C. S. T. Daughtry. 1990. Models for remote sensing of primary production at the regional and global scales. Proc. IGARSS'90 Symp. College Park, MD. 21-25 May 1990.

C. SYSTEMS RESEARCH

Acock, B. 1990. Effects of carbon dioxide on photosynthesis, plant growth, and other processes. In ASA Special Publication No. 53, Impact of Carbon Dioxide, Trace Gases, and Climate Change on Global Agriculture. pp 455-60.

Acock, B. 1990. Structuring agricultural research to deliver decision-support products. pp. 443-48. In: Proceedings of the Workshop on Computer-Integrated Agriculture. Agr. Res. Inst., Bethesda, MD.

Acock, B. 1991. Modeling canopy photosynthetic response to carbon dioxide, light interception, temperature, and leaf traits. In CSSA Special Publication Number 19, Modeling Crop Photosynthesis - From Biochemistry to Canopy.

Acock, B., M. C. Acock, and D. Pasternak. 1990. Interactions of CO₂ enrichment and temperature on carbohydrate production and accumulation in muskmelon leaves. J. Amer. Soc. Hort. Sci. 115:525-529.

Acock, B. and M. C. Acock. 1991. Potential for using long-term field research to develop and validate crop simulators. Agron. J. 83:46-61.

Acock, B. and J. F. Reynolds. 1990. Model structure and database development. pp. 169-179. In: R. K. Dixon, R. S. Meldahl, G. A. Ruark, and W. G. Warren, (eds). Process Modeling of Forest Growth Responses to Environmental Stress. Timber Press, Portland, Oregon.

Acock, B. and A. Trent. 1991. The soybean crop simulator, GLYCIM: documentation for the modular version 91. Miscellaneous Series Bulletin 145. Idaho Agricultural Experiment Station, University of Idaho, College of Agriculture.

Lestrade, J. P., B. Acock, and A. Trent. 1990. The effect of cloud layer plane albedo on global and diffuse insolation. Solar Energy 44:115-121.

Reddy, V. R. and D. N. Baker. 1990. Application of GOSSYM to analysis of the effects of weather on cotton yields. Agr. Sys. 32:83-95.

Reddy, V. R., D. N. Baker, and H. F. Hodges. 1991. The effects of temperature on cotton canopy growth, photosynthesis and respiration. Agron. J. 83:699-704.

Reddy, V. R., H. F. Hodges, and K. R. Reddy. 1990. Streamlining agricultural information from the scientist to the economic users through simulation models. Intl. Agric. Eng. Conf., Bangkok, Thailand, 3-6 Dec. 1990. pp. 1213-1222.

Reddy, V. R., K. R. Reddy, and D. N. Baker. 1991. Temperature effect on growth and development of cotton during the fruiting period. Agron. J. 83: 211-217.

Reddy, V. R., K. R. Reddy, H. F. Hodges, and D. N. Baker. 1990. The effect of temperature on growth, development, and photosynthesis of cotton during the fruiting period. pp. 97-110. In: Mechanisms of plant perception and response to environmental stimuli. British Society for Plant Growth Regulation, Monograph No. 20.

D. BIOCHEMICAL AND PHYSIOLOGICAL STUDIES

Agrawal, M., S. Agrawal, D. T. Krizek, G. F. Kramer, E. H. Lee, R. M. Mirecki, and R. A. Rowland. 1991. Physiological and morphological responses of snapbean plants to ozone stress as influenced by pretreatment with UV-B radiation. pp. 133-146. In: Y. Abrol et al. (eds.), Impact of Global Climatic Changes on Photosynthesis and Plant Productivity. Oxford & IBH Publ. Co., New Delhi, India.

Avni, A., R. A. Mehta, A. K. Mattoo, B. M. Greenberg, B. B. Chattopadhyay, D. Heller, and M. Edelman. 1991. Nucleotide sequence of the Spirodela oligorrhiza chloroplast psbA gene coding for the D1 (32kDa) photosystem II protein. Plant Mol. Biol. 17:919-921.

Bennett, J. H., E. H. Lee, and H. E. Heggestad. 1990. Inhibition of photosynthesis and leaf conductance interactions induced by SO₂, NO₂ and SO₂ + NO₂. Atmos. Environ. 24:557-562.

Britz, S. J. 1990. Photoperiodic and thermoperiodic regulation of assimilate partitioning in leaves of crop plants. pp. 853-866. In: D. K. Hayes, J. E. Pauly, and J. R. Reiter (eds.) Chronobiology: Its Role in Clinical Medicine, General Biology, and Agriculture, Part B., Alan R. Liss, Inc., New York, pp. 940.

Britz, S. J. 1990. Photoregulation of root:shoot ratio in soybean seedlings. Photochem. Photobiol. 52:151-159.

Britz, S. J. 1990. Regulation of photosynthate partitioning into starch in soybean leaves: response to natural daylight. Plant Physiol. 94:350-356.

Britz, S. J. 1990. Photomorphogenesis and photoassimilation in soybean and sorghum grown under broad spectrum or blue-deficient light sources. Plant Physiol. 94:448-454.

Britz, S. J. 1991. Setting the clocks that time photosynthate partitioning into starch stored soluble sugar in Sorghum: response to spectral quality. pp. 265-274. In: J. L. Bonnemain, S. Delrot, W. J. Lucas and J. Dainty (eds.) Recent Advances in Phloem Transport and Assimilate Compartmentation. Ouest Editions, Nantes, France.

Bunce, J. A. 1990. Short- and long-term inhibition of respiratory carbon dioxide efflux by elevated carbon dioxide. Ann. Bot. 65: 637-642.

Bunce, J. A. and F. Caulfield. 1991. Reduced respiratory carbon dioxide efflux during growth at elevated carbon dioxide in three herbaceous perennial species. Ann. Bot. 67:325-330.

De Veau, E. J., J. M. Robinson, R. D. Warmbrodt, and P. Van Berkum. 1990. Photosynthesis and photosynthate partitioning in N₂-fixing soybeans. Plant Physiol. 94:259-267.

Drake, B. G., P. W. Leadley, L. H. Ziska, and W. T. Pockman. 1989. Elevated atmospheric CO₂ reduces dark respiration in two halophytes. Plant Physiol. 89(4):40.

Foy, C. D., E. H. Lee, C. A. Coradetti, and G. J. Taylor. 1990. Organic acids related to differential aluminum tolerance in wheat (*Triticum aestivum*) cultivars. pp. 381-389. In: M. L. van Beusichem (ed.) Plant Nutrition-Physiology of Applications, Kluwer Academic Publishers, Boston, MA.

Ghirardi, M. L., S. K. Sopory, T. D. Elich, B. M. Greenberg, M. Edelman, and A. K. Mattoo. 1991. Light-induced degradation of the D1 photosystem II reaction center protein. In: G. S. Singhal and T. Ramasarma (eds.), Trends in Bioenergetics and Biotech. Proc., Today & Tomorrow's Printers and Publishers, New Delhi, India.

Heggestad, H. E. 1989. Air pollution injuries to pome fruits. pp. 218-224. In: P. R. Fridlund (ed.) Virus and virus-like diseases of pome fruits and simulated non-infectious disorders. No. SP-0003. Washington State Univ. Pullman, WA.

Heggestad, H. E. 1991. Origin of Bel W-3, Bel-C and Bel-B tobacco varieties and their use as indicators of ozone. Environ. Pollut. 74:264-291.

Heggestad, H. E., V. M. Lesser. 1990. Effects of ozone, sulfur dioxide, soil water deficit, and cultivar on yields of soybean. J. Env. Qual. 19:488-495.

Heggestad, H. E. and E. H. Lee. 1990. Soybean root distribution and top growth in response to ambient ozone and soil moisture stress when grown in soil columns in greenhouses. Environ. Pollut. 65:195-207.

Hogan, K. P., A. P. Smith, and L. H. Ziska. 1991. Potential effects of elevated CO₂ and changes in temperature on tropical plants. *Plant, Cell and Environ.* 14:763-778.

Kramer, G. F., E. H. Lee, C. L. Mulchi, and R. A. Rowland. 1991. Effect of increased ozone and CO₂ levels on field-grown soybeans. Polyamine titre. *Environ. Pollut.* 73:137-152.

Kramer, G. F., E. H. Lee, R. A. Rowland, and C. L. Mulchi. 1991. Effects of elevated CO₂ concentration on the polyamine levels of field-grown soybean at three O₃ regimes. *Environ. Pollut.* 73:137-152.

Kramer, G. F., H. A. Norman, D. T. Krizek, and R. M. Mirecki. 1991. Influence of UV-B radiation on polyamines, lipid peroxidation and membrane lipids in cucumber. *Phytochemistry* 30:2101-2108.

Krizek, D. T. 1989. The changing atmosphere: Implications for agriculture. *Proc. Agric. Res. Inst.* 37:21-35.

Krizek, D. T., R. M. Mirecki, R. P. Philbeck, and F. L. Booker. 1990. An improved lamp/filter system for use in UV enhancement studies under greenhouse and field conditions. Pp. 271-278. In: H. D. Payer, T. Pfirrmann & P. Mathy (eds.), *Air Pollution Research Report 26. Environmental Research With Plants in Closed Chambers*, Commission of the European Communities, Munich, Germany.

Lee, E. H. 1990. Soil, Water and Air. USDA Quarterly Report of Selected Research Project. p.9. 1 April - 30 June 1990.

Lee, E. H. 1991. Plant resistance mechanisms to air pollutants: Circadian rhythms in ascorbic acid production in response to ozone stress. *Chronobiology* 8:93-102.

Lee, E. H., R. A. Rowland, and C. L. Mulchi. 1990. Growth regulators serve as a research tool to study the mechanism of plant response to air pollution stimuli. *British Plant Growth Regulator Group, Monograph 20.* pp 127-137.

Pausch, R. C., S. J. Britz, and C. L. Mulchi. 1991. Growth and photosynthesis of soybean (Glycine max (L.) Merr.) in simulated vegetation shade: influence of the ratio of red to far-red radiation. *Plant, Cell Environ.* 14:647-656.

Robinson, J. M. 1990. Spinach leaf chloroplast NO₂⁻ photoreduction is stimulated by fructose 1, 6 biphosphate and/or triose phosphates. *Current Topics in Plant Biochem. and Physiol.* 9:416.

Robinson, J. M. 1990. Nitrite photoreduction in vivo is inhibited by oxygen. *Plant Physiol.* 92:862-865.

Roeb, G. and S. J. Britz. 1991. Short-term fluctuations in the transport of assimilates to the ear of wheat measured with steady-state ^{14}C - CO_2 -labeling of the the flag leaf. J. Expt. Bot. 42:469-475.

Slaughter, L. H., C. L. Mulchi, E. H. Lee, and K. Tuthill. 1989. Chronic ozone stress effects on yield and grain quality of soft red winter wheat. Crop Sci. 29:1251-1255.

Teramura, A. H., J. S. Sullivan, and L. H. Ziska. 1990. The interaction of elevated UV-B radiation and CO_2 on productivity and photosynthesis in rice, wheat and soybean. Plant Physiol. 4(2):470-475.

Teramura, A. H., L. H. Ziska, and A. Ester Sztein. 1991. Changes in growth and photosynthetic capacity of rice with increased UV-B radiation. Physiol. Plant. 83:373-380.-

Vigil, E. L., E. F. Erbe, and H. D. Fahimi. 1990. Application of stereological and image analysis to examination of organelle and membrane structure. pp. 585-588. In: EMAG-MICRO 89 Vol. 2: Biological. H. Y Elder and P. J. Goodhew (eds.) Institute of Physics Conference Ser. No. 98. IOP Publishing Ltd. Bristol, England.

Vigil, E. L. and H. D. Fahimi. 1990. A quantitative freeze-fracture investigation of intramembranous particles during the transformation of glyoxysomes to peroxisomes in watermelon cotyledons. Eur. J. Cell Biol. 51:1-8.

Vigil, E. L., T. K. Fang, D. P. Thibodéaux, C. Conkerton, O. Hinojosa, and J. Evans. 1991. Progress report on the relationship between fiber quality and seed quality as affected by drought stress. pp. 859-860. In: Proc. Cotton Beltwide Production Conf.

Whitaker, B. D., E. H. Lee, and R. A. Rowland. 1990. EDU and ozone protection: Foliar glycerolipds and steryl lipids in snapbean exposed to O_3 . Physiol. Plant. 80:286-293.

Wu, Y., E. H. Lee, and E. M. Barrows. 1989. Effects of SO_2 on Mexican bean beetle. Acta Ecologica Sinica. 9:336-340.

Wu, Y., E. H. Lee, and E. M. Barrows. 1990. The effect of ozone on feeding and growth of the Mexican bean beetle larvae. Acta Entomologica Sinica. 33: 161-165.

Ziska, L. H. 1990. Effects of salinity on photosynthesis. McGraw-Hill Yearbook of Science and Technology, 10th ed., 1991, McGraw-Hill, New York, pp 328-331.

Ziska, L. H. and A. H. Teramura. 1991. The interaction of elevated CO_2 and UV-B radiation on photosynthesis and productivity in rice (Oryza sativa). Plant Physiol. 91(4):23.

Ziska, L. H., S. J. Chamberlain, and B. G. Drake. 1989. The effect of elevated CO₂ during growth in the field on photosynthesis and respiration of a C₃ and C₄ salt marsh species. Bull. Ecol. Soc. Amer. 70(2):21.

Ziska, L. H., T. M. DeJong, G. J. Hoffman, and R. M. Mead. 1991. Sodium and chloride distribution in salt-stressed Prunus salicina, a deciduous tree species. Tree Physiol. 8:47-57.

Ziska, L. H., B. G. Drake, and S. Chamberlain. 1989. Long-term effects of elevated CO₂ on photosynthesis and dark respiration of a C₃ and C₄ species in the field. Plant Physiol. 89(4):163

Ziska, L. H., B. G. Drake, and S. Chamberlain. 1990. Long-term photosynthetic response in single leaves of a C₃ and C₄ salt marsh species grown at elevated atmospheric CO₂ in situ. Oecologia. 83:469-472.

Ziska, L. H., K. P. Hogan, A. P. Smith, and B. G. Drake. 1991. Growth and photosynthetic response of nine tropical plant species with long-term exposure to elevated carbon dioxide. Oecologia. 86:383-389.

Ziska, L. H., H. E. Reed, J. H. Sullivan, and A. H. Teramura. 1990. Stomatal and biochemical limitations to photosynthesis in soybean cultivars differing in sensitivity to UV-B radiation. Bull. Ecol. Soc. Amer. 71(2):379-380.

Ziska, L. H., J. R. Seemann, and T. M. DeJong. 1990. Determination of salinity induced limitations to the photosynthetic process in Prunus salicina, a deciduous tree species. Plant Physiol. 93:864-870.

SECTION 3

PUBLISHED ABSTRACTS

1989 - 1991

1991 ANNUAL REPORT ON GLOBAL CLIMATE CHANGE RESEARCH AT BARC

A. HYDROLOGIC STUDIES (Abstracts)

Kustas, W. P., Perry, E. M., and Doraiswamy, P. C. 1991. MONSOON '90 - A remote sensing feasibility study of the energy and water balance of a semiarid basin. Preprint IAF-91-160 of the 42nd Congress of the International Astronautical Federation, 5-11 Oct. Montrael, Canada. p 6. (Abstract)

Kustas, W. P., Schmugge, T. J., Jackson, T. J., Goodrich, D. C., Bach, L. B., Moran, M. S., Stannard, D. I., Hipps, L. E. and Blanford, J. H. 1991. Effects of localized precipitation on basin-scale hydrologic cycle. EOS. 72:117. (Abstract)

Rango, A. 1991. Worldwide testing of the Snowmelt Runoff Model with applications for predicting the effects of climate change. Nordic Hydrology. 22, p 10. (Abstract)

Rango, A. 1991. Worldwide testing of SRM with applications for predicting the hydrologic effects of climate change (in extreme years). American Geophysical Union. EOS. p. 189. (Abstract)

B. STUDIES ON REMOTE SENSING (Abstracts)

Andresen, J. A., K. P. Gallo, and C. S. T. Daughtry. 1989. Linking modeled crop responses and satellite derived vegetation indices. Agron. Abstr. 81:11.

Daughtry, C. S. T. and K. P. Gallo. 1990. Spectral estimates of absorbed radiation and phytomass in plant canopies. p. 18. In: Remote Sensing for Agriculture. Proc. Beltsville Symp. Greenbelt, MD. 16-18 May 1990. (Abstract)

Daughtry, C. S. T., W. P. Kustas, M. S. Moran, and R. D. Jackson. 1989. Spatial variability of net radiation and soil heat flux in agricultural fields. Agron. Abstr. 81:12-13.

Daughtry, C. S. T., M. A. Weltz, E. M. Perry, and W. P. Dulaney. 1991. Direct and indirect measurements of leaf area index. Agron. Abstr. 83:15.

Gallo, K. P. and C. S. T. Daughtry. 1989. The use of remotely sensed data in computation of crop radiation utilization. Agron. Abstr. 81:15.

Gallo, K. P. and C. S. T. Daughtry. 1990. The use of remotely sensed data in computation of crop radiation utilization. p. 39. In: Remote Sensing for Agriculture. Proc. Beltsville Symp. Greenbelt, MD. 16-18 May 1990.

Goward, S. N., S. D. Prince, C. S. T. Daughtry, and D. G. Dye. 1990. Macroscale models of primary production using remote sensing inputs. p. 20. In Remote Sensing for Agriculture. Proc. Beltsville Symp. Greenbelt, MD. 16-18 May 1990. (Abstract)

Perry, E. M., C. S. T. Daughtry, and W. P. Kustas. 1990. Relation between instantaneous and daily total absorbed photosynthetically active radiation over soybeans. p. 42. In: Remote Sensing for Agriculture. Proc. Beltsville Symp. Greenbelt, MD. 16-18 May 1990. (Abstract)

Prince, S. D., S. N. Goward, D. G. Dye, and C. S. T. Daughtry. 1990. Models for remote sensing of primary production at the regional and global scales. Proc. IGARSS'90 Symp. College Park, MD. 21-25 May 1990. (Abstract)

Ranson, K. J., J. R. Irons, and C. S. T. Daughtry. 1990. Canopy albedo from bidirectional reflectance. p. 24. In Remote Sensing for Agriculture. Proc. Beltsville Symp. Greenbelt, MD. 16-18 May 1990. (Abstract)

C. SYSTEMS RESEARCH (Abstracts)

Acock, B. and V. R. Reddy. 1991. Modeling cotton responses to temperature. ASA Meetings, Denver, Colorado, 27 Oct.- 1 Nov. 1991. (Abstract)

Hodges, H. F. and V. R. Reddy, and K. R. Reddy. 1990. Effects of temperature on flowering and fruit retention in cotton. ASA Meetings, San Antonio, TX, 21-26 Oct. 1990. (Abstract)

Lemmon H., and N. Chuk. 1992. Salient physiological and mathematical features of a new cotton model. Proc. Beltwide Cotton Prod. Meetings, Knoxville, TN, 10 Jan. 1992. (Abstract)

Reddy, K. R., H. F. Hodges, and V. R. Reddy. 1990. The effects of CO₂ on cotton growth and photosynthesis during fruiting. ASA Meetings, San Antonio, TX, 21-26 Oct. 1990. (Abstract)

Reddy, V. R., K. R. Reddy, and B. Acock. 1991. Carbon dioxide enrichment and temperature effects on growth and photosynthesis of cotton. ASA Meetings, Denver, Colorado, 27 Oct.- 1 Nov. 1991. (Abstract)

Reddy, V. R., K. R. Reddy, and B. Acock. 1991. The effects of carbon dioxide enrichment and temperature on growth and photosynthesis of cotton. Proc. Beltwide Cotton Prod. Res. Conf., San Antonio, TX, 8-12 Jan. 1991. (Abstract)

Reddy, V. R., K. R. Reddy, and M. C. Acock. 1991. The effects of aerial carbon dioxide enrichment and temperature on root growth in cotton. Proc. Beltwide Cotton Prod. Res. Conf., San Antonio, TX, 8-12 Jan. 1991. (Abstract)

Reddy, V. R., K. R. Reddy, and D. N. Baker. 1990. Responses of cotton canopies to temperature. In T. Cotton Nelson (ed.) Proc. Beltwide Cotton Prod. Res. Conf., Las Vegas, NV, 9-14 Jan. 1990. (Abstract)

Reddy, V. R., K. R. Reddy, and H. F. Hodges. 1990. Developing suitable database for modeling crop response to weather. p. 13. Workshop on crop simulation, Biological Systems Simulation Group, 27-29 March 1990, Lincoln, NE. (Abstract)

Trent, A., B. Acock, V. R. Reddy, and M. C. Acock. 1991. Validation of soybean model GLYCIM with different cultivars and climatic conditions. ASA Meetings, Denver, Colorado, 27 Oct.- 1 Nov. 1991. (Abstract)

C. BIOCHEMICAL AND PHYSIOLOGICAL STUDIES (Abstracts)

Adamse, P. and S. J. Britz. 1991. Role of photosynthesis in amelioration of UV-B damage. Plant Physiol. 96(S):114. (Abstract)

Britz, S.J. 1990. Setting the clocks that time photosynthate partitioning into starch and stored sugar in Sorghum: response to spectral quality. Proc. Intl. Conf. Cognac, France, 19-24 Aug. 1990. (Abstract)

Bunce, J. A. and F. Caulfield. 1990. Reduced respiration at elevated atmospheric carbon dioxide in three herbaceous perennials. Bull. Ecol. Soc. Amer. 71(S):108. (Abstract)

Bunce, J. A. and D. R. Lee. 1991. Light, temperature and nutrients as factors in photosynthetic adjustment to elevated carbon dioxide. Plant Physiol. 96(S):132. (Abstract)

De Veau, E. J. and J. M. Robinson. 1989. Effects of N₂ fixation and supplemental nitrogen on photosynthesis and photosynthate partitioning in soybeans. Plant Physiol. 89(S):18. (Abstract).

De Veau, E. J. and J. M. Robinson. 1990. Influences of N₂ fixation on net photosynthesis and chlorophyll content in soybean leaves. Plant Physiol. 93(S):125. (Abstract).

De Veau, E. J., J. M. Robinson, and R. D. Warmbrodt. 1990. Influences of N₂ fixation on partitioning of fixed carbon between carbohydrate and protein in soybean leaves. Plant Physiol. 93(S):125. (Abstract).

Drake, B. G., P. W. Leadley, L. H. Ziska, and W. T. Pockman. 1989. Elevated atmospheric CO₂ reduces dark respiration in two halophytes. Plant Physiol. 89(4):40.

- Elich, T.D., M. Edelman and A.K. Mattoo. 1991. Identification and characterization of a phosphorylated form of the 32kDa-D1 reaction center protein from *Spirodella oligorrhiza*. International Meeting on Regulation of Chloroplast Biogenesis, 28 July - 3 Aug. 1991, Crete, Greece. p. 22. (Abstract)
- Kramer, G. F., D. T. Krizek and R. M. Mirecki. 1991. Influence of visible and UV-B radiation on polyamine levels in soybean. *Plant Physiol.* 96(S):114. (Abstract)
- Krizek, D. T., M. Agrawal, S. B. Agrawal, G. F. Kramer, E. H. Lee, R. M. Mirecki, and R. A. Rowland. 1991. Morphological and physiological responses of cucumber to inverse day/night temperature and ozone pollution. *HortScience* 26(6):174. (Abstract)
- Mattoo, A.K., T.D. Elich and M. Edelman.. 1991. Does redox-regulated phosphorylation target the D1 protein of photosystem II for degradation? Beltsville Symposium XVI Morphogenesis in Plants: College Park, MD., 22-27 Sept. 1991. p. S26. (Abstract),
- Norman, H. A., D. T. Krizek, and R. M. Mirecki. 1989. Lipid compositional changes during low-temperature pre-conditioning against SO₂ in coleus. *Plant Physiol.* 89(S):28. (Abstract)
- Robinson, J. M. 1989. Oxygen inhibits NO₂⁻ photoreduction in plastid and cell isolates. *Plant Physiol.* 89(S):473. (Abstrac).
- Robinson, J. M. 1989. Chloroplast NO₂⁻ photoreduction is enhanced by CO₂ fixation. *Plant Physiol.* 89(S):1193. (Abstract).
- Robinson, J. M. 1990. Spinach leaf chloroplast NO₂⁻ photoreduction is stimulated by fructose-1,6-bisphosphate and/or triose phosphates. *Current Topics in Plant Biochemistry and Physiology* 9:416. (Abstract).
- Robinson, J. M. and E. J. De Veau. 1990. Selecting soybean leaves which yield mesophyll cell isolates displaying maximal rates of CO₂ and NO₂⁻ photoassimilations. *Plant Physiol.* 93(S):125. (Abstract).
- Vigil, E. L. and T. K. Fang. 1990. Stage specific effect of severe drought on precocious germination of developing cotton (*Gossypium hirsutum*, L. Var. M-8) embryos. *Plant Physiol.* 93(S):13. (Abstract)
- Ziska L, H., B. G. Drake, and S. Chamberlain. 1989. Long-term effects of elevated CO₂ on photosynthesis and dark respiration of a C₃ and C₄ species in the field. *Plant Physiol.* 89(4):163. (Abstract)

Ziska, L. H, S. J. Chamberlain, and B. G. Drake. 1989. The effect of elevated CO₂ during growth in the field on photosynthesis and respiration of a C3 and C4 salt marsh species. Bull. Ecol. Soc. Amer. 70(2):21. (Abstract)

Ziska, L. H. and A. H. Teramura. 1991. The interaction of elevated CO₂ and UV-B radiation on photosynthesis and productivity in rice (Oryza sativa). Plant Physiol. 91(4):23. (Abstract)

